

ARTICLE

Open Access

The H⁺-pyrophosphatase *IbVP1* regulates carbon flux to influence the starch metabolism and yield of sweet potato

Weijuan Fan^{1,2}, Yandi Zhang^{1,3}, Yinliang Wu^{1,3}, Wenzhi Zhou¹, Jun Yang^{1,2}, Ling Yuan⁴, Peng Zhang^{1,3} and Hongxia Wang¹

Abstract

Storage roots of sweet potato are important sink organs for photoassimilates and energy, and carbohydrate metabolism in storage roots affects yield and starch production. Our previous study showed that sweet potato H⁺-pyrophosphatase (*IbVP1*) plays a vital role in mitigating iron deficiency and positively controls fibrous root growth. However, its roles in regulating starch production in storage roots have not been investigated. In this study, we found that *IbVP1* overexpression in sweet potato improved the photosynthesis ability of and sucrose content in source leaves and increased both the starch content in and total yield of sink tissues. Using ¹³C-labeled sucrose feeding, we determined that *IbVP1* overexpression promotes phloem loading and sucrose long-distance transport and enhances Pi-use efficiency. In sweet potato plants overexpressing *IbVP1*, the expression levels of starch biosynthesis pathway genes, especially *AGPase* and *GBSSI*, were upregulated, leading to changes in the structure, composition, and physicochemical properties of stored starch. Our study shows that the *IbVP1* gene plays an important role in regulating starch metabolism in sweet potato. Application of the *VP1* gene in genetic engineering of sweet potato cultivars may allow the improvement of starch production and yield under stress or nutrient-limited conditions.

Introduction

Sweet potato (*Ipomoea batatas* (L.) Lam.) is a multi-functional starchy root crop species that is a primary resource for food, feed, and bioindustrial applications. The annual global production of sweet potato is approximately 92 million tons of fresh storage roots (FAOSTAT 2018), which is crucial for food security, malnutrition and poverty alleviation, and bioindustrial development^{1,2}. With increasing demand for sweet potato production, storage roots need to be improved,

especially in terms of yield and quality, both of which are positively correlated with the acquisition and partitioning of organic carbon^{3,4}. In addition to a strong photosynthesis ability of the leaves, an efficient system of sucrose transport from source to sink via stem phloem can ensure sufficient access to carbohydrates^{5–7}. Sucrose transporters (SUTs) are responsible for sucrose loading into the complex comprising phloem sieve elements and companion cells. Overexpression of *SUT* genes has been demonstrated to improve plant productivity by effective partitioning of carbohydrates in sinks in rice, pea, and wheat^{8–10}. In sweet potato, overexpression of the H⁺-translocating inorganic pyrophosphatase (H⁺-PPase) gene *IbVP1* increases plasma membrane H⁺-ATPase hydrolytic activities, *SUT1* expression, and root growth¹¹. Nevertheless, how *IbVP1* regulates starch metabolism in sweet potato storage roots has not yet been determined.

Correspondence: Peng Zhang (zhangpeng@cemps.ac.cn) or Hongxia Wang (hxwang@cemps.ac.cn)

¹National Key Laboratory of Plant Molecular Genetics, CAS Center for Excellence in Molecular Plant Sciences, Chinese Academy of Sciences, Shanghai 200032, China

²Shanghai Key Laboratory of Plant Functional Genomics and Resources, Shanghai Chenshan Plant Science Research Center, Chinese Academy of Sciences, Shanghai 201602, China

Full list of author information is available at the end of the article

© The Author(s) 2021

 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

H⁺-PPase involves in plant metabolism by hydrolyzing pyrophosphate to form sodium and/or proton gradients that transport ions across the membrane^{12,13}. Under abiotic stress conditions, H⁺-PPase gene expression and enzyme activity increase^{14–17}. H⁺-PPase affects the transport and accumulation of auxin and sugar during root and shoot growth^{18–22}. These phenomena lead to enhanced root and shoot biomass; increased yields of grain, fiber, or fruit; and improved photosynthesis ability, nutrient and water uptake, and CO₂ fixation (especially under nutrient-deficient conditions, e.g., low NO₃⁻ and phosphate (Pi))^{9,23–25}. Overexpression of H⁺-PPase in *Arabidopsis* results in the upregulated expression of the sucrose proton symporter *SUC1*, suggesting the involvement of H⁺-PPase in the transport of sucrose into the phloem^{18,19}.

In a previous study, we showed that overexpressing *IbVP1* in sweet potato improves plant growth and development through altered carbohydrate metabolism in the leaves, transport of polar auxin from source to sink, and rhizosphere acidification to facilitate nutrient acquisition¹¹. However, the underlying molecular mechanism for the function of *IbVP1* in storage root starch metabolism remains unclear. In this study, we showed that overexpression of *IbVP1* resulted in an increase in starch accumulation in, storage root numbers of and yield of sweet potato. Transgenic sweet potato plants exhibited increased phloem loading and long-distance transportation of sucrose through the acquisition and utilization of activated Pi, thereby increasing the starch content in and yield of the storage roots. Moreover, the starch physicochemical properties, including their composition and structure, were altered due to the increased transcript level of *granule-bound starch synthase I (GBSSI)*. These findings advance our mechanistic understanding of the functions of *IbVP1* in the starch biosynthesis process and provide a new target for adding value to root crops by improving starch content and quality.

Materials and methods

Transgenic sweet potato lines

Sweet potato cv. Taizhong 6, one of the edible yellow-fleshed cultivars grown in China, was used as a donor parent for generating transgenic plants. *IbVP1*-overexpressing transgenic lines (IA lines) were produced as described by Fan et al.¹¹. In early May 2014–2017, wild-type (WT) plants and transgenic lines were planted in a field at the Wushe Plantation for Transgenic Crops, Shanghai, China (31°13948.0099°N, 121°28912.0099°E).

Starch isolation from sweet potato storage roots

The storage roots of IA lines and WT plants were harvested after the plants had grown in the field for 5 months. Freshly harvested storage roots of the sweet

potato plants were washed, peeled, cut into small slices, suspended in distilled water, and blended with a commercial scale blender. The starchy liquid was passed through a 100 μm sieve, and the starch granules were allowed to sediment. The supernatants were decanted and replaced four times with distilled water. The final supernatants were discarded, and the samples were dried at 40 °C for two days²⁶.

Analysis of sugar and starch contents

Storage roots of WT and IA lines growing in the field for 5 months were harvested, ground into powder, and then heated at 80 °C for approximately 48 h to obtain a constant dry weight (DW). The dried samples (30 mg) were used to determine the sugar and starch contents, as described by Knutson et al.²⁷. The starch content was measured by the use of starch kit (Megazyme International Ireland Limited, Wicklow, Ireland). After sugar extraction, the supernatant was centrifuged and transferred into a new glass tube for high-performance liquid chromatography (HPLC) analysis for the identification of the sugar fractions. The different fractions of sugar were identified on the basis of their retention time via known standards. The sugar concentrations were subsequently determined with a standard curve.

Amylose content measurements

The amylose contents in the starch samples from 5-month-old field-grown WT and IA plants were measured by the colorimetric method by following a previously described protocol²⁸. Pure amylose (Type III, Sigma A0512, St. Louis, MO) and amylopectin (Sigma 10118) of potato starch were purchased and used to make known dilutions to establish standard curves.

Characterization of starch thermal properties

The thermal properties of starch from 5-month-old field-grown WT and IA plants were assessed by the differential scanning calorimeter method (DSC Q2000; TA Instrument, Ltd., Crawley, UK). All the samples were prepared and kept in aluminum sealed pans for 2 h, after which scanning was performed in the temperature range from 30 to 95 °C. An empty, sealed aluminum pan was used as a reference. The measurements were recorded in triplicate for each individual sample.

Pasting properties of starch

The pasting properties of starch from 5-month-old field-grown WT and transgenic plants were measured using a Rapid Visco-Analyzer (RVA) (RVA-4 series, Newport Scientific, Warriewood, Australia). A starch suspension (5% wt/vol) was produced by mixing in distilled water, and starch viscosity parameters were recorded at various temperatures as described by Zhou et al.²⁶.

Particle size distribution assays

The distribution of granule size of starch from 5-month-old field-grown WT and IA lines were determined by using a Master-sizer (2000 Laser Diffraction Instrument Malvern Instruments, Ltd., Worcestershire, UK) in wet-cell mode as described by Zhou et al.²⁶ and Blazek et al.²⁹.

Morphology of starch granules

The granular morphology of the starch from 5-month-old field-grown plants was observed via scanning electron microscopy (SEM). Starch samples were dispersed in distilled water and then spread onto double-sided adhesive tape, air dried, and coated with gold powder. The samples were observed via SEM (JSM6360IV, JEOL, Tokyo, Japan) and imaged. For transmission electron microscopy (TEM), slices of freshly prepared storage roots (1 mm³) were glazed with 1,3-diformal propane. Afterward, ultrathin sectioning was performed, and the samples were examined via TEM (Hitachi H7650, Tokyo, Japan).

Measurements of amylopectin chain length distribution (CLD)

Amylopectin CLD was measured by using the method of Nishi et al.³⁰. First, the isolated sweet potato starch was debranched enzymatically for quantification of the chain-length distribution. In brief, 5 mg of sweet potato starch sample was digested with isoamylase enzymes (Sigma; 15284). The debranched starch samples were then used to quantify the amylopectin content using high-performance anion-exchange chromatography with pulsed amperometric detection (HPAEC-PAD).

Freeze–thaw stability assays

The stability of starch subjected to freeze–thaw cycles from 5-month-old field-grown plants was evaluated by a previously described method²⁸. Briefly, 5% starch paste (wt/vol) was liquified by boiling for 20 min. The samples were cooled, after which 0.5 mL of each sample was transformed into new 1.5 mL Eppendorf tubes. The weight of the starch in each sample tube was measured. All the samples were stored at –70 °C overnight and then put into a water bath to thaw for 60 min at 22 °C. Six replications from each starch tube were taken and centrifuged at 8000 *xg* for 10 min at 18 °C. The supernatant was separated, and the remaining paste was weighed. The obtained value (represented as the percent of the initial sample; wt/wt) was considered a measure of syneresis. The tubes were subsequently stored at –70 °C for further freeze–thaw cycles and measurements.

Phloem loading measurements

The phloem exudate assay was performed on plants grown for 3 months in a greenhouse as described previously³¹, with some modifications. Stems with three

unfolded leaves were cut and soaked immediately in a 15 mM EDTA solution. The stems were then recut to pieces 1.5 to 2 cm in length in 15 mM EDTA solution buffer to collect the phloem exudate. After the samples incubated for 12 h in the dark at 25 °C, phloem exudates were concentrated for sucrose measurements via HPLC.

The photoassimilates were labeled with stable ¹³C₂ isotopes to investigate the diffusion rate of photoassimilates into the phloem, as described by Gui et al.³² and Koubaa et al.³³. The first unfolded leaf was positioned in a gas-sealed transparent glass chamber filled with ¹³C₂. After 12 h of photosynthesis, treated plants were kept in the dark, and leaf samples were collected for sucrose isolation and measurements. Following derivatization, analysis was performed by gas chromatography–mass spectrometry (GC–MS). The proportions of ¹³C isotope mass spectral fragments were determined to measure the amounts of ¹³C-labeled sugars.

Measurements of leaf area

Leaves from sweet potato plants that had been growing for 3 months (six different pots) were sampled to measure the leaf area. The leaves were imaged, and the leaf area (mm²) was measured using WSeen software. At least six replicates for each leaf were recorded for the transgenic lines and WT.

Photosynthesis measurements

Net photosynthesis was measured on the third leaf from the top of plants grown for three months in the field, using a LI-6400 photosynthesis system (LI-6400 Inc., Lincoln, NE, USA). All measurements were carried out at 9:00 to 10:00 a.m. following the manufacturer's instructions.

Sucrose phosphate synthase (SPS) activity assays

For SPS analysis, the first unfolded leaves of 3-month-old plants were frozen in liquid nitrogen, after which total proteins were extracted, desalted using a spin column, and immediately assayed. SPS activity was determined using a sucrose phosphate synthase assay kit (Solarbio BC0600) according to the manufacturer's specifications.

Activity assays of storage root enzymes

Vacuole H⁺-PPase activity was determined using the method of Fan et al.¹¹.

The enzyme activity of starch biosynthesis-related enzymes (AGPase, SS, GBSS, and SBE) was measured in tuberous roots of the AI and WT plants by following the methods described by Nakamura et al.³⁴. One unit of enzyme (SS, GBSS, and AGPase) activity was defined as the production of 1 nmol of ADP per min at 30 °C, while 1 unit of SBE enzyme activity was described as the amount of enzyme needed to increase the spectrophotometric absorbance by one unit at 540 nm in 1 min. The activities of α-amylase and β-amylase in the storage

roots of sweet potato plants were also measured. A Cer-alpha kit (Megazyme International Ireland, Bray Business Park, Bray, Co., Wicklow, Ireland.) was used following the manufacturer's protocol, with the appropriate dilutions.

Expression of starch biosynthesis enzymes

The mRNA levels of starch biosynthesis-related genes in the storage roots of IA and WT plants were measured by quantitative real-time reverse transcription polymerase chain reaction (qRT-PCR). Gene-specific primers were designed (Table S3) to analyze the expression levels using SYBR Green PCR Master Mix (Bio-Rad) in a Bio-Rad CFX96 thermocycler. Amplification was performed under the following conditions: 95 °C for 1 min, followed by 40 cycles of 95 °C for 15 s and 60 °C for 30 s. The relative expression levels of the genes were calculated using the sweet potato *Actin* gene as an internal control.

Pyrophosphate determination

For the quantification of pyrophosphate (PPi), roots from 5-month-old field-grown plants were collected and immediately frozen in liquid nitrogen to avoid PPi hydrolysis. The frozen roots were homogenized in chilled 80% ethanol, vortexed for 10 min, heated at 80 °C for 30 min, and then centrifuged at 20,000 × *g* for 5 min. The precipitate was subsequently resuspended in chilled 80% ethanol and centrifuged. The supernatants were collected and dried with a centrifugal evaporator. The dried extract was resuspended in water, and water-soluble compounds were extracted. PPi was measured using a pyrophosphate assay kit (Sigma MAK168-1KT) according to the manufacturer's specifications. All glassware was pretreated overnight with 0.1 M HCl to remove residual Pi.

Determination of Pi in sweet potato tissue

The Pi content in roots and shoots of 5-month-old field-grown plants was determined by digesting the tissues in H₂SO₄-H₂O₂. After 2 mL of H₂SO₄ was added to the samples followed by thorough shaking (including the blowing of air through a pipette), 1 mL of H₂O₂ was added two times via multiple small drops. After the reaction was completed, the sample was placed in an electric heater (nitrogen blowing at 120 °C for 15 min) until the solid disappeared into an essentially transparent solution. The digested solutions were subsequently diluted with ddH₂O. Aliquots of diluted solutions (0.5 mL) were added to 4 mL of a solution consisting of 0.5% (w/v) ammonium molybdate, 0.6 M H₂SO₄, and 2% (w/v) ascorbic acid. The absorbance was read at 660 nm, and the Pi content was measured as milligrams of Pi per gram of fresh weight.

Statistical analysis

At least three biological replicates were used for each experiment. The data are reported as the means ± SDs, with

three replicates for each individual experiment. Significant differences were identified for each individual treatment using Student's *t* test via SigmaPlot 10.0 (Systat Software, San Jose, CA). The differences among the treatments were statistically significant when *P* was <0.05 or 0.01.

Results


IbVP1-overexpressing sweet potato lines presented increased yield and higher phosphorus accumulation

We generated stable *IbVP1* transgenic sweet potato lines and characterized the impacts of *IbVP1* overexpression on shoot and fibrous root growth¹¹. However, the storage roots of the *IbVP1* transgenic lines have not been thoroughly studied. To understand the impact of *IbVP1* overexpression on storage root yield, mature storage roots from 5-month-old *IbVP1*-overexpressing sweet potato plants (IA lines) were harvested from the field. Compared with the WT plants, all IA lines exhibited higher yields and increased numbers of storage roots (Fig. 1a, b, Fig. S1a-c). Under field conditions, the yield of the IA lines ranged from 7.47 to 8.53 kg/plant, which is 20–39% higher than the WT yield 6.23 kg/plant (Fig. 1b). IA plants produced a higher number of storage roots 6.3 ± 1.5, 6.3 ± 1.2, and 5.7 ± 1.2 than did the WT plants (3.7 ± 1.2) (Fig. S1c).

IbVP1 hydrolyzes pyrophosphate to Pi. The V-H⁺-PPase hydrolytic activities of IA and WT roots were determined. As shown in Fig. 1c, the H⁺-PPase activity in the IA plants was higher than that in the WT plants. Moreover, the PPi concentrations decreased, but the Pi concentrations increased more in the three IA lines compared to the WT (Fig. 1d–e). Furthermore, the Pi concentrations of the IA leaves were 41% higher than those of the WT leaves (Fig. 1f), indicating that the IA lines have more Pi available in both leaf and root cells. Pi is an indispensable macro-nutrient essential for plant growth and development³⁵. The higher concentrations of Pi in the IA lines occurred simultaneously with improved root growth.

IbVP1 overexpression increases carbohydrate export and translocation from source to sink

To assess the source photoassimilates, leaf area and average photosynthesis rates were measured. Compared with the WT, IA7 and IA8 had larger leaf areas (Fig. 2a). The net photosynthesis of the IA leaves was higher than that of WT leaves (Fig. 2b). Sucrose levels significantly increased in the leaves of the IA lines compared with the WT (Fig. S2a). However, the starch contents in the IA leaves were lower than those in the WT leaves (Fig. S2b). Therefore, the ratio of sucrose/starch in the leaves of the IA plants was significantly higher than that of the WT (Fig. S2c). Furthermore, the SPS activities in the IA leaves increased by 1.2- to 1.8-fold, which likely enhanced sugar biosynthesis in the IA leaves compared to WT leaves. To investigate the relationship between photoassimilate



translocation and increased yields in the IA plants, we determined the amount of sucrose exported from the leaves. Sucrose levels in phloem exudates collected from the IA lines were significantly higher than those from the WT (Fig. 2c). Additionally, photoassimilate loading from leaves into the phloem was also recorded by using the ^{13}C stable isotopic labeling method combined with GC-MS. The sucrose contents in the leaves of the IA lines were higher than those in the WT (Fig. 2d). Consistent with these findings, more ^{13}C -labeled sucrose was detected in the fibrous roots of IA plants compared with the WT, indicating an increase in the rate of sucrose transport out of leaves of the IA lines (Fig. 2e). Taken together, these results clearly demonstrate that the transgenic IA lines are more efficient at exporting carbohydrates, mainly sucrose, from photosynthetic leaves, promoting the yield of storage roots via upregulation of *lbVP1* expression.

Storage root development is strongly associated with starch accumulation³⁶. To determine whether starch metabolism is affected by *lbVP1* expression in sweet potato, the levels of major carbohydrates (i.e., fructose, glucose, sucrose, and starch) in the storage roots of IA and WT plants were measured. The sucrose contents in the IA lines were 28–67% higher than those in the WT plants (Fig. 3a). Glucose and fructose were also higher in IA lines

than in the controls (Fig. 3a). The total starch and amylose contents in the IA lines were, respectively, 20–32% and 27–32% higher than those in WT, which were significantly different (Fig. 3b–c).

lbVP1 overexpression enhances starch biosynthesis ability

Since the *lbVP1*-overexpressing sweet potato plants showed an increased carbon flux during starch biosynthesis, we measured the expression of genes involved in starch biosynthesis. We found that the expression levels of the *AGPase*, *GBSSI*, *SSI*, *SBEI*, and *SBEII* genes were upregulated in the transgenic lines (Fig. 4a). In addition, the mRNA levels of the genes encoding α -amylase and β -amylase, two starch-degrading enzymes, were significantly decreased (Fig. 4a).

The corresponding activities of AGPase, GBSSI, SSI, and SBEs increased significantly in transgenic lines by 1.14- to 1.26-, 2.03- to 2.57-, 1.69- to 1.96-, and 1.16- to 1.32-fold, respectively, compared with those in the WT (Fig. 4b), whereas α -amylase and β -amylase activities significantly decreased in the IA lines (Fig. 4b). Taken together, these findings suggest that increases in the activities of AGPase, GBSSI, SSI, and SBEs lead to increased starch deposition in the tuberous roots of IA lines.




Fig. 2 Sucrose translocation in sweet potato. **a** Leaf area, **b** net photosynthesis rate, and **c** SPS activity were measured in wild-type (WT) and IA lines. **d** Sucrose contents in phloem exudates of WT and *lbVP1*-overexpressing transgenic (IA lines) sweet potato. **e–f** ¹³C-labeled sucrose in the leaves (**e**) and fibrous roots (**f**) of sweet potato plants after feeding photosynthetic leaves ¹³CO₂. The asterisks represent significant differences compared to WT at **P* < 0.05 or ***P* < 0.01 (Student’s *t* test)





Fig. 3 Contents of major carbohydrates in sweet potato storage roots. **a** Fructose, glucose, and sucrose. **b** Starch. **c** Amylose. WT wild type; IA4, IA7, and IA8 are three independent *lbVP1*-overexpressing transgenic lines. Five-month-old storage roots harvested from the field were used for analysis. The asterisks represent significant differences from WT at **P* < 0.05 or ***P* < 0.01 (Student’s *t* test)



***lbVP1* overexpression affects both granular starch size and morphology in sweet potato storage roots**


The starch extracted from the storage roots of IA lines showed a broad distribution of granular starch size, ranging from 5.2 to 58.9 μm in the IA lines compared to 5.2–51.9 μm in the WT, and the average granule sizes of the IA lines were higher than those of the WT (Fig. 5a). Furthermore, the representative granule diameters, Dx10, Dx50, and Dx90, in the IA lines were 11, 20.1, and 36.2 μm, respectively, which were significantly higher than the 10.3, 17.8, and 29.9 μm diameters, respectively, of granules from the WT storage roots (Fig. 5b).

SEM analysis did not reveal any significant differences among the starch granule shapes between the IA lines and WT. Different granule shapes (oval, round, or polygonal) within the starch were observed for both IA lines and WT (Fig. 5c). However, some cracks were observed in the starch isolated from the IA roots, which is consistent with that observed previously in high-amylose sweet potato starch, as high amylose content disrupts the structure of the starch semicrystalline structure^{26,37}. The cross-sections of sweet potato tuberous roots were examined under TEM, and WT starch granules revealed “zebra

stripe” patterns, whereas those from IA transgenic lines showed darker black stripes (Fig. 5d).

***lbVP1*-overexpressing sweet potato plants present altered starch composition and properties**

The pasting profiles of IA and WT starch were measured with RVA. As the temperature increased, granules of the WT starch swelled, with a rapid increase in viscosity; the granule structures of starch also became distorted, resulting in polymer dispersion, and finally, starch gel formed upon cooling. As shown in Fig. 6a, the characteristic curves of the IA starch were dramatically different from those of the WT starch, with the former exhibiting no discernible peak. We also measured the pasting property parameters, i.e., peak viscosity, final viscosity, breakdown, peak time, setback, and pasting temperature (Table S1). The viscosity peaks ranged from 1159 cP to 1290 cP for the IA starch; IA7 starch presented the lowest viscosity peak (1159 cP), which was still higher than that of the WT (1101 cP). The lowest breakdown values ranged from 175 cP (in IA7) to 252 cP (in IA8), and both were higher than the 114 cP of the WT (Table S1). The final viscosity was higher in the IA plants than the



WT. However, the pasting temperatures of the IA starch were lower than those of the control starch.


Clear differences in onset temperature (T_o), top melting temperature (T_p), end melting temperature (T_e), and enthalpy (ΔH) were observed in the IA starch compared with the WT starch. During the melting process, the T_o decreased slightly for all IA starch, ranging from 63.5 °C (in IA4) to 60.32 °C (in IA8), compared to the WT T_o of 64.77 °C, with a ΔH of 11.32 J g⁻¹. Notable decreases in T_p and T_e were also detected in the IA starch compared to the WT starch (Fig. 6b). To determine the freeze–thaw stability in sweet potato starch, the level of syneresis of 5% starch paste was measured by repeating several cycles of freezing/thawing. Our results showed that, compared with the WT starch, the IA starch displayed significantly higher syneresis, even after just one cycle, ranging from 36.13–50.96%, with a maximum syneresis of 60.8% (Fig. 6c).

Amylopectin CLD was measured in isoamylase-treated starch samples using HPAEC-PAD analysis. The degree of polymerization (DP 6–75) was determined and normalized based upon the peak area (Fig. 7). The patterns of the amylopectin peaks were quite consistent among all the tested samples despite their different levels. Notably, the first trough was observed at DP 8, while two peaks were observed at DP 11–13 and at approximately DP 43, and there was a slight hump at DP 18; these results are

consistent with previous findings^{26,38,39}. To verify the DP differences, the amylopectin glucan chain value of WT was deduced from the corresponding IA starch (Fig. 7b–d). The chain lengths of DP 10–14 and DP 34–75 were reduced, whereas the DP 15–23 chains increased in the IA lines. Comparative starch analysis was performed, and it was noted that, compared with the WT starch, the IA starch comprised fewer short chains (maximum, –0.1%) and slightly more long chains (maximum, 0.28%).

Discussion

Root crop productivity is largely determined by the ratio of acquisition and partitioning of organic carbon to sink organs. Previously, the roles of H⁺-PPases in many plants were reported to be associated with enhancing shoot and root growth by the efficient delivery of increased amounts of fixed carbon to sink tissue^{18–25}. Although starch is the major form of fixed carbon in root crops, the roles of *IbVP1* in the storage roots of root crops have not yet been described. In our study, we found that overexpression of *IbVP1* in sweet potato resulted in increased photosynthesis ability of and sucrose contents in source leaves and promoted phloem loading and sucrose transport to sink organs, which in turn increased the total starch content in and yield of storage roots (Figs. 1–3). Gene expression and the activities of starch biosynthesis enzymes were concurrently upregulated (Fig. 4). *IbVP1*-induced *AGPase* gene





expression is perhaps particularly critical for increased starch biosynthesis and total starch content because AGPase is the rate-limiting enzyme in starch metabolism^{40,41}. *IbVP1* overexpression increased storage root numbers of the transgenic plants compared to the WT plants, possibly due to increased source-to-sink sucrose loading and transport for starch biosynthesis, triggering the development of some fibrous roots into storage roots³⁶. Our results illustrate the important roles of *IbVP1* in increasing yield and increasing storage root numbers.

Sufficient carbohydrates from photosynthesis and high rates of phloem loading and sucrose transport are essential for the partitioning of photoassimilates into sink organs for plant production. Our data indicated that *IbVP1* enhances the net photosynthesis of leaves and influences sucrose translocation (Fig. 2) by assisting sucrose loading into

phloem, thus resulting in more efficient sucrose transport from source to sink organs. In addition, upregulated expression of the sucrose proton symporter *SUT1* in sweet potato leads to higher sucrose levels in storage roots, suggesting that improved sucrose transport from the leaves to the roots enhances carbon fixation in storage roots (Figs. 1 and 3)¹¹. Similarly, transgenic *Arabidopsis* lines with phloem companion cell-specific expression of *Arabidopsis* Vacuolar Pyrophosphatase (AVP) consistently have more hexose and starch in rosettes due to improved phloem transport accompanied by upregulated *Suc* gene expression¹⁷. Over-expression of the *AVP1* gene in *Arabidopsis* and barley results in an increase in phloem loading, transport, and unloading of sucrose into sink organs, which leads to biomass enhancement^{19,22,42}. Improved transport of sugars from source to sink in H^+ -PPase-overexpressing plants may explain the increased biomass production as well as increased fixed carbon in the root system^{21,25}. In our study, compared with the field-grown WT plants, the field-grown IA transgenic lines showed enhanced growth and vigor as well as increased yields of and total starch contents in the storage roots (Fig. 1). Sweet potato plants overexpressing *IbVP1* show enhanced levels of sucrose transport from source to sink, which likely affects auxin transport and increases rhizosphere acidification, leading to improved nutrient and water uptake¹⁹. Compared with the WT, the transgenic plants had 41 and 10% higher Pi concentrations in the leaves and roots, respectively, and the yields of storage roots of the IA lines were 20–39% higher than those of the WT in the field. In rice, overexpression of H^+ -PPase results in root systems that are more robust than those of the WT under both Pi-sufficient and Pi-deficient conditions¹⁷, which signifies the role of H^+ -PPase in optimizing Pi-use efficiency in plants^{17,25}.

In addition to the increased total starch content of the IA lines, the ratio of amylose to amylopectin changed, which likely affects the physicochemical and functional properties of starch^{43,44}. The amylose content peaked at 28.67% in the IA lines compared with 21.75% in the WT (Fig. 3c). In the IA lines, the expression of *GBSSI*, which is responsible for amylose biosynthesis, seems to be higher than the expression of *SBEI* and *SBEII*, which are responsible for amylopectin biosynthesis. This differential expression may explain the increase in the relative proportion of amylose. Furthermore, starch properties, including pasting properties and gelatinization, were altered in the IA lines (Figs. 5–6). We observed increased gelatinization temperature, lower breakdown, and higher setback values in starch from the IA lines compared with the WT. These values are similar to the high-amylose starch in the *SBE*-RNAi lines of sweet potato, which have higher proportions of long-chain amylopectin and larger granule sizes²⁶. High amylose starch is desirable in some food applications because it is highly resistant to digestion and has a lower



absorbance response^{43,44}, thereby reducing health problems such as obesity and diabetes^{45,46}.

In summary, sweet potato is an important root crop, and increasing yield and improving starch quality, especially increased storage root numbers and increased amylose contents, are difficult objectives to achieve through traditional breeding. Prior to this study, it was unclear how *IbVP1* functions in the storage roots of sweet potato. Our study showed that *IbVP1* plays a significant role in starch metabolism by altering the carbon metabolism flux to improve starch metabolism and increase yields of storage roots by influencing Pi metabolism. Enhanced phloem loading and long-distance transport of sucrose, mediated by *IbVP1*, increase yields and total starch content by mobilizing more photoassimilates from sources to sink organs (storage roots)

for sink biomass production through increased photosynthesis, enlarged leaf areas, improved SPS activity, and decreased starch levels in the leaves of *IbVP1*-overexpressing transgenic lines. Overexpression of *IbVP1* increased rhizosphere acidification by increasing plasma membrane H⁺-ATPase activity to promote nutrient acquisition and auxin transport, contributing to improved plant growth, especially under stress conditions¹¹. We suggest that *IbVP1* functions to enhance sucrose transport mediated by the sucrose proton symporter SUT1 and to increase starch biosynthesis ability by altering carbon metabolism and starch physicochemical properties. These outcomes lead to enhanced carbohydrate transport from source organs to sink organs, increasing both root development and starch content (Fig. 8). The substantial improvement in yield due to

increased tuber numbers and starch quality of the IA lines implies a practical approach to breeding sweet potato by manipulation of *IbVPI*.

Acknowledgements

This work was supported by grants from the National Key R&D Program of China (2018YFD1000700, 2018YFD1000705, 2019YFD1000701-2), the National Natural Science Foundation of China (31501356), the Special Funds for the Local Science and Technology Development of the Central Government (2020ZY0006), the Key Laboratory of Biology and Genetic Improvement of Tuber and Root Crops, Ministry of Agriculture and Rural Affairs, China (No. NYBSL201801), the National Key Technology Research and Development Program of China (2015BAD15B01), and the Natural Science Foundation of Shanghai (18ZR1413300).

Author details

¹National Key Laboratory of Plant Molecular Genetics, CAS Center for Excellence in Molecular Plant Sciences, Chinese Academy of Sciences, Shanghai 200032, China. ²Shanghai Key Laboratory of Plant Functional Genomics and Resources, Shanghai Chenshan Plant Science Research Center, Chinese Academy of Sciences, Shanghai 201602, China. ³University of Chinese Academy of Sciences, Beijing 100049, China. ⁴Department of Plant and Soil Sciences and Kentucky Tobacco Research and Development Center, University of Kentucky, Lexington, KY 40546, USA

Author contributions

W.F. performed most of the experiments, analyzed the data, and drafted the manuscript. Y.Z., Y.W., and W.Z. conducted part of the starch analysis. J.Y. provided helpful suggestions for the experiments. L.Y. analyzed the data and revised the manuscript. P.Z. and H.W. designed the study, analyzed the data, and revised the manuscript. All the authors have read and approved the final manuscript.

Conflict of interest

The authors declare that they have no conflict of interest.

Supplementary Information accompanies this paper at (<https://doi.org/10.1038/s41438-020-00454-2>).

Received: 9 July 2020 Revised: 19 November 2020 Accepted: 28 November 2020

Published online: 01 February 2021

References

- Bovell, B. A. C. Sweet potato: a review of its past, present, and future role in human nutrition. *Adv. Food Nutr. Res.* **52**, 1–59 (2007).
- Liu, Q. C., Liu, J., Zhang, P. & He, S. Z. Root and Tuber Crops. In *Encyclopedia of agriculture and food systems*. (ed. Van Alfen, N.) Vol. 5, 46–61 (Elsevier, 2014).
- Fedoroff, N. V. et al. Radically rethinking agriculture for the 21st century. *Science* **327**, 833–834 (2010).
- Godfray, H. C. J. et al. Food security: the challenge of feeding 9 billion people. *Science* **327**, 812–818 (2010).
- Ainsworth, E. A. & Bush, D. R. Carbohydrate export from the leaf: a highly regulated process and target to enhance photosynthesis and productivity. *Plant Physiol.* **155**, 64–69 (2011).
- Braun, D. M. Plant science. SWEET! The pathway is complete. *Science* **335**, 173–174 (2012).
- Chen, J. H. et al. Nuclear-encoded synthesis of the D1 subunit of photosystem II increases photosynthetic efficiency and crop yield. *Nat. Plants* **6**, 570–580 (2020).
- Rosche, E. et al. Seed-specific overexpression of a potato sucrose transporter increases sucrose uptake and growth rates of developing pea cotyledons. *Plant J.* **30**, 165–175 (2002).
- Weichert, N. et al. Increasing sucrose uptake capacity of wheat grains stimulates storage protein synthesis. *Plant Physiol.* **152**, 698–710 (2010).
- Saalbach, L. et al. Increased grain yield and micronutrient concentration in transgenic winter wheat by ectopic expression of a barley sucrose transporter. *J. Cereal Sci.* **60**, 75–81 (2014).
- Fan, W. J. et al. H⁺-pyrophosphatase *IbVPI* promotes efficient iron use in sweet potato [*Ipomoea batatas* (L.) Lam.]. *Plant Biotechnol. J.* **15**, 698–712 (2017).
- Luoto, H. H., Baykov, A. A., Lahti, R. & Malinen, A. M. Membrane integral pyrophosphatase subfamily capable of translocating both Na⁺ and H⁺. *Proc. Natl Acad. Sci. USA.* **110**, 1255–1260 (2013).
- Baykov, A. A., Cooperman, B. S., Goldman, A. & Lahti, R. Cytoplasmic inorganic pyrophosphatase. *Prog. Mol. Subcell. Biol.* **23**, 127–150 (1999).
- Arif, A., Zafar, Y., Arif, M. & Blumwald, E. Improved growth, drought tolerance, and ultrastructural evidence of increased turgidity in tobacco plants over-expressing *Arabidopsis* vacuolar pyrophosphatase (*AVP1*). *Mol. Biotechnol.* **54**, 379–392 (2013).
- Maeshima, M. Vacuolar H⁺-pyrophosphatase. *Biochim. Biophys. Acta* **1465**, 37–51 (2000).
- Paez, V. J. et al. Enhanced proton translocating pyrophosphatase activity improves nitrogen use efficiency in romaine lettuce. *Plant Physiol.* **161**, 1557–1569 (2013).
- Yang, H. et al. Enhanced phosphorus nutrition in monocots and dicots overexpressing a phosphorus-responsive type I H⁺-pyrophosphatase. *Plant Biotechnol. J.* **5**, 735–745 (2007).
- Gaxiola, R. A., Sanchez, C. A., Paez-Valencia, J., Ayre, B. G. & Elser, J. J. Genetic manipulation of a “vacuolar” H⁺-PPase: from salt tolerance to yield enhancement under phosphorus-deficient soils. *Plant Physiol.* **159**, 3–11 (2012).
- Khadiolkar, A. S. et al. Constitutive and companion cell-specific overexpression of *AVP1*, encoding a proton-pumping pyrophosphatase, enhances biomass accumulation, phloem loading, and long-distance transport. *Plant Physiol.* **170**, 401–414 (2016).
- Li, J. S. et al. *Arabidopsis* H⁺-PPase *AVP1* regulates auxin mediated organ development. *Science* **310**, 121–125 (2005).
- Pasapula, V. et al. Expression of an *Arabidopsis* vacuolar H⁺-pyrophosphatase gene (*AVP1*) in cotton improves drought and salt tolerance and increases fiber yield in the field conditions. *Plant Biotechnol. J.* **9**, 88–99 (2011).
- Pizzio, G. A. et al. *Arabidopsis* type I proton-pumping pyrophosphatase expresses strongly in phloem, where it is required for pyrophosphate metabolism and photosynthate partitioning. *Plant Physiol.* **167**, 1541–1553 (2015).
- Lv, S. L. et al. H⁺-pyrophosphatase from *Salicornia europaea* confers tolerance to simultaneously occurring salt stress and nitrogen deficiency in *Arabidopsis* and wheat. *Plant Cell Environ.* **38**, 2433–2449 (2015).
- Gaxiola, R. A. & Hirschi, K. D. Moving on up: H⁺-PPase mediated crop improvement. *Trends Biotechnol.* **34**, 347–349 (2016).
- Yang, H. B. et al. Over-expression of the *Arabidopsis* proton-pyrophosphatase *AVP1* enhances transplant survival, root mass, and fruit development under limiting phosphorus conditions. *J. Exp. Bot.* **65**, 3045–3053 (2014).
- Zhou, W. Z. et al. Impact of amylose content on starch physicochemical properties in transgenic sweet potato. *Carbohydr Polym.* **122**, 417–427 (2015).
- Knutson, C. A. & Grove, M. J. Rapid method for estimation of amylose in maize starches. *Cereal Chem.* **71**, 469–471 (1994).
- Zhou, W. Z. et al. Alpha-glucan, water dikinase 1 affects starch metabolism and storage root growth in cassava (*Manihot esculenta* Crantz). *Sci. Rep.* **7**, 9863 (2017).
- Blazek, J. & Copeland, L. Pasting and swelling properties of wheat flour and starch in relation to amylose content. *Carbohydr Polym.* **71**, 380–387 (2008).
- Nishi, A., Nakamura, Y., Tanaka, N. & Satoh, H. Biochemical and genetic analysis of the effects of amylose-exender mutation in rice endosperm. *Plant Physiol.* **127**, 459–472 (2001).
- Eom, J. S. et al. Impaired function of the tonoplast localized sucrose transporter in rice, *OssUT2*, limits the transport of vacuolar reserve sucrose and affects plant growth. *Plant Physiol.* **157**, 109–119 (2011).
- Gui, J. S., Liu, C., Shen, J. H. & Li, L. G. Grain setting defect1, encoding a remorin protein, affects the grain setting in rice through regulating plasmodesmal conductance. *Plant Physiol.* **166**, 1463–1478 (2014).
- Koubaa, M., Mghaieth, S., Thomasset, B. & Roscher, A. Gas chromatography–mass spectrometry analysis of ¹³C labeling in sugars for metabolic flux analysis. *Anal. Biochem.* **425**, 183–188 (2012).
- Nakamura, Y., Yuki, K., Park, S. Y. & Ohya, T. Carbohydrate metabolism in the developing endosperm of rice grains. *Plant Cell Physiol.* **30**, 833–839 (1989).

35. Chen, Y. F., Wang, Y. & Wu, W. H. Membrane transporters for nitrogen, phosphate and potassium uptake in plants. *J. Integr. Plant Biol.* **50**, 835–848 (2008).
36. Wang, H. X. et al. Altered phenylpropanoid metabolism in the Maize Lc-expressed sweet potato (*Ipomoea batatas*) affects storage root development. *Sci. Rep.* **6**, 18645 (2016).
37. Fulton, D. C. et al. Role of granule-bound starch synthase in determination of amylopectin structure and starch granule morphology in potato. *J. Biol. Chem.* **277**, 10834–10841 (2002).
38. Rolland, S. A. et al. Molecular and supra-molecular structure of waxy starches developed from cassava (*Manihot esculenta* Crantz). *Carbohydr Polym.* **92**, 1451–1462 (2013).
39. Kitahara, K. et al. Physicochemical properties of amylose-free and high-amylose starches from transgenic sweet potatoes modified by RNA interference. *Carbohydr Polym.* **69**, 233–240 (2007).
40. Dong, M. Y., Fan, X. W. & Li, Y. Z. Cassava *AGPase* genes and their encoded proteins are different from those of other plants. *Planta* **250**, 1621–1635 (2019).
41. Na, G., Aryal, N., Fathihi, A., Kang, J. L. & Lu, C. F. Seed-specific suppression of ADP-glucose pyrophosphorylase in *Camelina sativa* increases seed size and weight. *Biotechnol. Biofuels* **11**, 330 (2018).
42. Schilling, R. K. et al. Expression of the *Arabidopsis* vacuolar H⁺-pyrophosphatase gene (*AVP1*) improves the shoot biomass of transgenic barley and increases grain yield in a saline field. *Plant Biotechnol. J.* **12**, 378–386 (2013).
43. Granfeldt, Y., Drews, A. & Björck, I. Arepas made from high amylose corn flour produce favorably low glucose and insulin responses in healthy humans. *J. Nutr.* **125**, 459–465 (1995).
44. Regina, A. et al. High amylose wheat generated by RNA interference improves indices of large bowel health of rats. *Proc. Natl Acad. Sci.* **103**, 3546–3551 (2006).
45. Frost, G. et al. Glycaemic index as a determinant of serum HDL- cholesterol concentration. *Lancet* **535**, 1045–1048 (1999).
46. Salmeron, J. et al. Dietary fiber, glycemic load, and risk of non-insulin-dependent diabetes mellitus in women. *J. Am. Med Assoc.* **277**, 472–477 (1997).